



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

During the same period there were reported 21 cases of smallpox, with 2 deaths.

*Statistics for the month of August, 1904—Plague and smallpox.*

Maximum temperature, 27.5° C.; minimum, 22° C.; average, 24.69° C. Average relative humidity, 76.3; rainy days, 5; rainfall, 21 mm.

Total deaths 522, as follows: Beriberi, 29; cancer and other malignant tumors, 7; debility, congenital, 11; debility, senile, 24; diseases of circulatory system, 56; diseases of digestive system, 75; diseases of nervous system, 40; diseases of respiratory system, 29; diseases of urinary organs, 15; erysipelas, 2; malarial fevers, 18; plague, 56; smallpox, 2; syphilis, 4; pulmonary tuberculosis, 68; tuberculosis, other, 5; whooping cough, 1; violent deaths, not suicides, 4; stillborn, 41; other causes, 35.

During the month there were 50 cases of smallpox, 53 cases of confirmed plague, and 34 cases of suspected plague.

*Corrected statistics of plague in Bahia from occurrence of the first case to October 11, 1904.*

JULY.

Number of cases bacteriologically verified as plague.....	27
Deaths during the month from same.....	9
Cases suspected as plague encountered dead by health officers, not examined bacteriologically, but considered as plague.....	8
Cases which had symptoms of plague, but bacteriologically did not prove to be such and not so considered.....	6
Total cases considered as plague.....	35
Total deaths considered as plague.....	17

AUGUST.

Number of cases bacteriologically verified as plague.....	53
Deaths during month from verified cases.....	23
Cases suspected as plague encountered dead by health officers, not examined bacteriologically, but considered as plague.....	34
Cases which had symptoms of plague, but bacteriologically did not prove to be such and not so considered.....	8
Total cases considered as plague.....	87
Total deaths considered as plague.....	57

SEPTEMBER.

Number of cases bacteriologically verified as plague.....	17
Deaths during the month from verified cases.....	10
Cases suspected as plague, encountered dead by health officers, not examined bacteriologically, but considered as plague.....	12
Cases which had symptoms of plague but bacteriologically did not prove to be such and not so considered.....	5
Total cases considered as plague.....	29
Total deaths considered as plague.....	22

OCTOBER.

Number of cases bacteriologically verified as plague.....	4
Deaths during month from verified cases.....	2
Cases suspected as plague, encountered dead by health officers, not examined bacteriologically, but considered as plague.....	2
Total cases considered as plague.....	6
Total deaths considered as plague.....	4

The number of new cases seems to be diminishing and is confined to the locations where there have heretofore been cases.